

Towards a Complete Calculation of $\gamma\gamma \rightarrow 4f^{*\dagger\ddagger}$

E. Boos^{1,2§}, T. Ohl^{2¶}

¹Moscow State University, Russia

²Darmstadt University of Technology, Germany

IKDA 97/36, November 1997

Abstract

We present a general classification of all four fermion final states in $\gamma\gamma$ collisions and a calculation of cross sections below the W^+W^- -threshold.

The planned e^+e^- Linear Collider will provide the opportunity to study $\gamma\gamma$ -collisions with energies of several hundred GeV in the center of mass system. Of immediate interest will be gauge and Higgs boson production. However, neither the Higgs, nor the W or the Z will be directly observed and we typically have to study four fermion production processes, for which the resonant diagrams $\gamma\gamma \rightarrow VV \rightarrow 4f$ do *not* form gauge invariant subsets and calculations of $\gamma\gamma \rightarrow 4f$ beyond on-shell gauge boson production are required for precision studies.

As a step towards the construction of a complete event generator for $\gamma\gamma \rightarrow 4f$, we have introduced [1] a classification of all four fermion final states in $\gamma\gamma$ -collisions and the corresponding Feynman diagrams. Fig. 1 shows all six topologies appearing in the Feynman diagrams for $\gamma\gamma \rightarrow 4f$. The flavors of the final state fermions determine which of W , Z , γ , or g contribute for the lines labelled such. In a three generation standard model, there are 98 possible four fermion final states that fall into only eight different classes. They are listed in table 1 with the number of diagrams for each topology. The charged current classes (CCn) are gauge invariant completions of W^-

*Contribution to the proceedings of "The International Europhysics Conference on High-Energy Physics", 19-26 August 1997, Jerusalem, Israel.

[†]Supported by Deutsche Forschungsgemeinschaft.

[‡]Presented by E.B., supported by the HEP97 Organizing Committee and Russian Ministry of Science.

[§]e-mail: boos@ifh.de

[¶]e-mail: Thorsten.Ohl@Physik.TH-Darmstadt.de




Figure 1: The six topologies for $\gamma\gamma \rightarrow 4f$ in unitary gauge (see [1] for R_ξ -gauges).

Class		Q	T	D	S	B	M	Σ	$\mathcal{O}(\alpha_S)$	e. g.
$CC13$	l	1	4	2	0	4	2	13		$e^- \bar{\nu}_e \mu^+ \nu_\mu$
$CC21$	sl	1	6	2	2	6	4	21		$e^- \bar{\nu}_e u \bar{d}$
$CC31$	h	1	8	2	4	8	8	31		$u \bar{d} \bar{c} s$
$NC06$	l/sl	0	0	0	2	4	0	6		$\nu_e \bar{\nu}_e \mu^- \mu^+$
$NC20$	l/h	0	0	0	4	8	8	20	+10	$e^- e^+ e^- e^+$
$NC40$	l/sl/h	0	0	0	8	16	16	40	+20	$e^- e^+ \mu^- \mu^+$
$mix19$	l	1	4	2	2	8	2	19		$e^- e^+ \nu_e \bar{\nu}_e$
$mix71$	h	1	8	2	12	24	24	71	+20	$u \bar{u} d \bar{d}$

Table 1: The eight classes of diagrams in $\gamma\gamma \rightarrow 4f$.




Figure 2: Integrated cross sections for charged current final states below the W^+W^- -threshold (full circles: full calculation, open circles: $\gamma\gamma \rightarrow WW^*$).

pair production and, unlike $e^+e^- \rightarrow 4f$, there are *no* final states without multi-peripheral diagrams. The neutral current classes (NCn) do not contain any cubic or quartic gauge couplings. Finally, the mixed classes (*mixn*) correspond to final states that are invariant under the transformation from “charge exchange” to “charge retention” form and vice versa. For each class, one typical final state is given (see [1] for complete lists). The column $\mathcal{O}(\alpha_S)$ shows the number of one gluon exchange diagrams to be added in the neutral current classes for purely hadronic final states.

References

- [1] E. Boos and T. Ohl, Phys. Lett. **B407** (1997) 161.